Copyright © 2018 by Academic Publishing House Researcher s.r.o.



Published in the Slovak Republic European Journal of Medicine Has been issued since 2013.

E-ISSN: 2310-3434 2018, 6(1): 20-42

DOI: 10.13187/ejm.2018.1.20

www.ejournal5.com



VITA Intense – Product with Negative Oxidation-reduction Potential (ORP) as Important Quality for Antioxidant and Inhibition Growth of Tumor Cells Effects. Anti Aging Effects

Ignat Ignatov a,*, Yuliana Pesheva a

^a Scientific Research Center of Medical Biophysics, Bulgaria

Abstract

We studied the mathematical model of interaction with water VITA intense of LavaVitae company (Austria). In this report are submitted data about the interaction of VITA intense with water, obtained by non-equilibrium (NES) and differential-equilibrium energy spectrum (DNES) of water. The average energy (ΔE_{H=0}) of hydrogen H...O-bonds among individual molecules H₂O after treatment of VITA intense with water measured by NES- and DNES-methods is ΔE=-0.0136±0.0011 eV for VITA intense. This result suggests the restructuring of ΔE_{H.,, O} values among H₂O molecules with a statistically reliable increase of local extremums in DNES-spectra. The research is performed for VITA intense, with study of pH and oxidative reduction potential (ORP). There is review of the effects of the chemical composition of VITA intense – antiinflammatory, antioxidant etc. With methods NES we show the following effects - relaxing effect of nervous system, anti-inflammatory and inhibition of tumor cells. As results of these effects VITA intense has anti aging influence. The base of this influence is anti-inflammatory effect. This article deals with the review of the basic biophysical-biochemical and biological processes underlying the VITA intense. The author is studying their physical-chemical properties and biophysical and biological effects on human organism. Additionally, by using IR, NES, and DNES methods are investigated various samples of water from Bulgarian water springs: the melt water from Glacier Rosenlaui, Swiss Alps, as well as the human blood serum of people with excellent health and cancer patients between 50 and 70 years old. Other experiments were performed on a 1 % (v/v) solution of VITA intense in deionized water. As an estimation factor in NES and DNES was measured the values of the average energy of hydrogen bonds (ΔE_{H...O}) among H₂O molecules in water samples, as well as a local extremums in the NES and DNES-spectra of various samples of water and the human blood serum at E = -0.1387 eV and $\lambda = 8.95 \mu m$. For a group of people in critical condition of life and patients with malignant tumors the greatest values of local extremums in IR-, DNESspectra were shifted to lower energies relative to the control healthy group. Further we applied this method for calculation of percent distribution of H₂O molecules in all studied water samples according to energies of hydrogen bonds ranged from (-0.08 to -0.1387 eV). It was shown that mountain water is among the most important factors for human longevity and human health. The variety of ions (K+, Na+, Ca2+, Mg2+, Mn2+, Fe2+, Fe3+, Zn2+, SO42-, Cl-, HCO3-, CO32-), the chemical-physical parameters (pH, electroconductivity) and the decreased content of deuterium in studied water samples renders beneficial effects of these types of water on human health. We are applying the conclusions for the effects of mountain water on human health and longevity as base

* Corresponding author

E-mail addresses: mbioph@abv.bg (I. Ignatov)

-

for the conclusion on the effects of VITA intense. The base is also the results with methods NES and DNES on human serum and expected effects from VITA intense.

VITA intense has negative oxidative reduction potential (ORP) and this is special property of this product for antioxidant effects again free radicals.

Keywords: VITA intense, anti-inflammatory, antioxidant, anti-aging, mathematical model, NES, DNES, negative oxidative reduction potential (ORP).

1. Introduction

Water is the main substance of life. The human body of an adult person is composed from 50 to 55 % of water. With aging, the percentage of water in the human body decreases. Hence, the factor of water quality and its amount in organism is an essential factor for the research (Pocock et al., 1981; Howard, Hopps, 1986). Water is present in the composition of the physiological fluids in the body and plays an important role as an inner environment in which the vital biochemical processes involving enzymes and nutrients take place. Water also is the main factor for metabolic processes and aging (Ignatov, 2012). Earlier studies conducted by us have demonstrated the role of water, its structure, the isotopic composition and physical-chemical properties (pH, temperature) on the growth and proliferation of prokaryotes and eukaryotes in water with different isotopic content (Ignatov, Mosin, 2012; Ignatov, Mosin, 2013). These factors, the structure and composition of water are of great importance in many biophysical studies. The peculiarities of the chemical structure of the H₂O molecule and weak bonds caused by electrostatic forces and donor-acceptor interaction between hydrogen and oxygen atoms in H₂O molecules create favorable conditions for formation of directed intermolecular hydrogen bonds (O-H...O) with neighboring H₂O molecules, binding them into complex intermolecular associates which composition represented by general formula (H₂O)_n, where n can vary from 3 to 50 (Keutsch & Saykally, 2011). The hydrogen bond is a form of association between the electronegative O-atom and a H-atom, covalently bound to another electronegative O-atom, is of vital importance in the chemistry of intermolecular interactions, based on weak electrostatic forces and donor-acceptor interactions with chargetransfer (Pauling, 1960). It results from interaction between electron-deficient H-atom of one H₂O molecule (hydrogen donor) and unshared electron pair of an electronegative O-atom (hydrogen acceptor) on the neighboring H₂O molecule.

The product of LavaVitae VITA intense is combining of herbs from Alps, Aloe Vera, Green tea, Edelweiss, Aronia, Vitamins and Magnesium, Selenium. The research is with methods NES and DNES. There is research of ORP and pH and there are executing the conclusions from electrochemically activated waters – anolyte and catholyte for anti-inflammatory effects (Ignatov et al., 2014).

The aim of this research is to show the usefully of VITA intense on the base of the following results and conclusions. The research are from various samples of water from Bulgarian water springs: the melt water from Glacier Rosenlaui, Swiss Alps, as well as human blood serum of people with excellent health and cancer patients between 50 and 70 years old. In frames of this research on the water quality were investigated 415 people living in the municipalities of Teteven, Yablanitza. Ugarchin, Lukovit, Lovech district; Dolni Dabnik, Pleven district, Kuklen, Plovdiv district (Bulgaria), where is lived the most of long lived people and their siblings, were studied.

The authors also performed the research of 1% (v/v) solution of VITA intense on the distribution of H_2O molecules according to the energies of hydrogen bonds, as well as studies of the NES and DNES spectrum and the biophysical effect of this type of water on human body.

2. Materials and Methods

2.1. NES and DNES Spectral Analysis

The device for DNES spectral analysis was made by A. Antonov on an optical principle. For this was used a hermetic camera for evaporation of water drops under stable temperature (+22–24 °C) conditions. The water drops were placed on a water-proof transparent pad, which consists of thin maylar folio and a glass plate. The light was monochromatic with filter for yellow color with wavelength at $\lambda = 580\pm7$ nm. The device measures the angle of evaporation of water drops from 72.3° to 0°. The DNES-spectrum was measured in the range of -0.08– -0.1387 eV or $\lambda = 8.9-13.8$ µm using a specially designed computer program. The main estimation criterion in these studies was the average energy ($\Delta E_{H...0}$) of hydrogen O...H-bonds among H₂O molecules in water

samples and human blood serum.

2.2. Product of LavaVitae - VITA intense

The product LavaVitae VITA intense of 1 liter is including – Natural and high quality plant extracts from the Swiss Alps, Aronia, Edelweiss, Aloe Vera, Vitamin C (180.0 mg), Vitamin E (15.0 mg), Vitamin A (17.0 mg), Vitamin B₁ (1.5 mg), Vitamin B₂ (1.7 mg), Vitamin B₃ (17.0 mg), Vitamins B₅ (10.0 mg), Vitamin B₆ (4.2 mg), Vitamins B₃ (17.0 mg), Watersoluble B Vitamin (17.0 mg) Vitamin B₉, Vitamin M, Vitamin B₁₁, (0.4 mg), Vitamin B₉, (7.4 μ g), Vitamin D₃, (5.0 μ g), Magnesium (62.5 mg), Selenium (25 μ g).

2.3. Studying the Bulgarian Long Living People and Centenarians

Interviews have been conducted with 415 Bulgarian centenarians and long living people and their siblings. Their heredity, body weight, health status, tobacco consumption, physical activity, attitude towards life has been analyzed. With using DNES method was performed a spectral analysis of 15 mountain water springs located in municipalities Teteven and Kuklen (Bulgaria). The composition of water samples was studied in the laboratory of "Eurotest Control" (Bulgaria). Statistics methods were attributed to the National Statistical Institute of Bulgaria.

2.4. Studying the *Human B*lood Serum

1 % (v/v) solution of human blood serum was studied with the methods of IR-spectroscopy, non-equilibrium (NES) and differential non-equilibrium (DNES) spectral analysis. The specimens were provided by Kalinka Naneva (Municipal Hospital, Bulgaria). Two groups of people between the ages of 50 to 70 were tested. The first group (control group) consisted of people in good clinical health. The second group included people in critical health or suffering from malignant diseases.

2.5. IR-spectroscopy

IR-spectra were registered on Brucker Vertex ("Brucker", Germany) IR spectrometer (a spectral range: average IR – 370–7800 cm⁻¹; visible – 2500–8000 cm⁻¹; the permission – 0,5 cm⁻¹; the accuracy of wave number – 0,1 cm⁻¹ on 2000 cm⁻¹) and on Thermo Nicolet Avatar 360 Fourier-transform IR.

2.6. Statistical Processing of *Experimental D*ata

Statistical processing of experimental data was performed using the statistical package STATISTISA 6.0 using the Student's t- criterion (at p < 0.05).

3. Results and Discussions

3.1. Applications of VITA intense for Human Health. The information is from the company LavaVitae

Product features: LavaVitae Company -VITA intense product

- Two bottles (2x500 ml) covers up the need for a monthly (~ 33 ml per a day);
- A synergy of essential vitamins and minerals with natural high-quality plant extracts and aromas from the Swiss Alps;
 - Highest bioavailability in liquid form;
- Without GMO. Free of animal ingredients and allergens according to regulation (EU) No. 1169/2011;
 - Supports a healthy diet and an active lifestyle.

The Table 1 shows the chemical composition of VITA intense

Table 1. The chemical composition of VITA intense

| Nutritional Information | Quantity in g per Serving (33 ml) | Quantity in g per 100 ml | |
|--------------------------------|--------------------------------------|-----------------------------|--|
| Calorific value | 18,53 kcal / 77,66 kJ | 56,14 kcal / 235,33 kJ | |
| Fat | < 0,1 | < 0,1 | |
| saturated fatty acids | <0,05 | <0,1 | |
| Carbohydrates | 4,5 | 13,67 | |
| of which sugar | 4,43 | 13,43 | |
| Protein | < 0,05 | < 0,1 | |
| Balla substances | 0,3 | 0,9 | |
| Salt | < 0,05 | < 0,1 | |

| Nutrient | Quantity in mg per Serving (33 ml) |
|-------------------------|------------------------------------|
| (Vitamins and Minerals) | |
| Vitamin A | 0,750 |
| Vitamin B1 | 1,5 |
| Vitamin B2 | 1,7 |
| Niacin | 17,0 |
| Pantothenic acid | 10,0 |
| Vitamin B ₆ | 4,2 |
| Biotin | 0,150 |
| Folic acid | 0,400 |
| Vitamin B ₁₂ | 0,0075 |
| Vitamin C | 180,0 |
| Vitamin D ₃ | 0,005 |
| Vitamin E | 15,0 |
| Magnesium | 62,5 |
| Selenium | 0.0025 |

| Another Nutrients | Quantity in mg per Serving (33 ml) | Quantity in mg per 100 ml |
|---------------------------|------------------------------------|------------------------------|
| Organic Aronia juice | 1,980 | 6.000 |
| Organic Aloe Vera extract | 17,5 | 53,0 |
| Edelweiss extract | 17,5 | 53,0 |
| Green tea extract | 6,9 | 21,0 |

Vitamins and Minerals:

Vitamin A

Improve eyesight and the immune system. Contributes to the maintenance of normal skin.

Vitamin B Complex

It strengthens nerves and ensures continuous supply of power. In the bustle of everyday life and times of increased willingness to perform the essential water-soluble vitamins support the metabolism and increase the stress tolerance. Thus counteract against shiftlessness and fatigue.

Vitamin B₁

Contributes to the normal function of the nervous system, psyche and heart.

Vitamin B₂

Contributes to the normal production of red blood cells and metabolism of iron.

Protect cells against oxidative stress.

Vitamin B₃ or Niacin

Helps maintaining normal skin.

Vitamin B₅ or Pantothenic acid

Contributes to the normal synthesis and metabolism of steroid hormones, Vitamin D and some neurotransmitters.

Vitamin B₆

Contributes to the normal metabolism of proteins and glycogen, to the normal production of red blood cells and regulating hormone balance.

Water-soluble B Vitamin other names: Vitamin H, Coenzyme R, Biotin

Helps maintaining normal skin, hair.

Vitamin B₉, Vitamin M, Vitamin B₁₁

Contributes to the normal synthesis of amino acid, formation of blood and plays an important role in cell division. Contributes to the normal growth of maternal tissues during pregnancy.

Vitamin R.

Contributes to the normal formation of red blood cells and plays an important role in cell division.

Vitamin C

Anti oxidant. Supports formation of collagen for skin, blood vessels and bone and aids in wound healing. Protects cells from oxidative stress. Supports the immune system. Increases the excretion of heavy metals via the kidneys and improves iron absorption.

Vitamin B₃

Contributes to the normal function of the muscles and maintaining normal bone and teeth. Contributes to the normal utilization of calcium and phosphorus and contributes to the normal calcium level in blood.

Vitamin E

Protect cells against oxidative stress and contributes to a young, tight and healthy skin. Increases the amount of collagen in the skin. Supports wound healing through the support of cell division.

Magnesium

Contributes to the normal muscle function and maintaining to normal bone and teeth. Supporting protein synthesis, regulates electrolyte balance and plays an important role in cell division.

Selenium

Contributes to the normal function of the immune system and the thyroid. Protect cells against oxidative stress.

Helps maintaining normal hair and nails as well as for normal formation of sperm. Natural and high quality plant extracts from the Swiss Alps. Extracts from plants provide an effective and long lasting protection against free radicals. For this reason Vita Intense contains a number of effective and health promoting extracts which are carefully matched to achieve an optimum synergistic effect with maximum bioavailability.

Aronia (Aronia melanocarpa)

The Aronia (Fig. 1) or "Black chokeberry" originates from North America is now increasingly cultivated in Middle Europe, including Switzerland.



Fig. 1. Aronia

The shell of the Aronia is naturally rich in vitamins (such as Vitamin A, C, E, K and the entire group of the B Vitamins), minerals and trace elements (such as calcium, magnesium, potassium, zinc and iron). The Aronia shows record antioxidants levels for protecting cells against oxidative stress (Zheng, Wang, 2003). But the most interesting active ingredient is hidden in the group of polyphenols, whose main representatives are the flavonoids and anthocyanins. The Aronia contains an unsurpassed amount of flavonoids and the highest proportion of anthocyanins compared to other berries (Table 2) (Zheng, Wang 2003).

Table 2. Composition of different berries (Zheng, Wang 2003)

| | – | |
|-----------------------------|--------------------------------|----------------------------|
| ORAC, Anthocyan- und Phenol | gehalt der Beeren im Vergleich | (nach Zheng und Wang 2003) |

| Beerensorte | ORAC (!mol Trolox Equivalent/g FG | Anthocyane (mg Cyanidin- 3-Glukoside/g FG) | Phenole (mg Gallussäureequivalent/gFG) |
|--------------|--------------------------------------|---|---|
| Blaubeere | 28,9 | 1,20 | 4,12 |
| Cranberries | 18,5 | 0,32 | 3,15 |
| Preiselbeere | 38,1 | 0,45 | 6,52 |
| Apfelbeere | 160,2 | 4,28 | 25,56 |

Figure 2 shows the results of anti oxidative parameter of Aronia (mmol/l) for TEAC.




Fig. 2. Oxidative parameter of Aronia (mmol/l) for TEAC

The Aronia has antioxidant (Olas et al., 2008), anti-inflammatory (Ohgami et al. 2005; Borisova et al., 1994), anti oxidative stress (Valcheva-Kuzmanova et al., 2005) antibacterial and antiviral effect (Kokotkiewicz et al., 2010). Further these antioxidants show positive effects on blood sugar (Ryszawa et al., 2005) and of blood lipid levels (Valcheva-Kuzmanova et al., 2006). They protect the heritage ~antimutagenic effect: (Gasiorowski et al., 1997; Dimova et al., 1997), the cardiovascular system (Ryszawa et al., 2006), the gastrointestinal tract (Valcheva-Kuzmanova et al., 2005; Matsumoto et al., 2004) and the liver (Borisova et al., 2004). While flavonoids improve blood circulation and the cardiovascular system (via relaxation of the vessel walls), anthocyanins reduce platelet formation (Ryszawa et al., 2006). Both flavonoids show unsurpassed antiviral, antimicrobial, antiallergic and anticarcinogenic effects (Bermudez-Soto et al., 2006; Lala, Marlik, 2006; Zhao et al., 2004; Marlik, 2003, Zhao et al., 2003).

Green tea (Camelia sinensis)

Green tea (Fig. 3) has been shown to preventative against many diseases. Green tea has been proven for everyday primary care, prevention of mild illnesses or weakened constitution, improving fitness and metabolism or for setting an ideal weight. (Lambert, Yang, 2003).



Fig. 3. Green tea

The extract of Green tea (*Camelia sinensisis*) naturally rich in many vitamins, minerals and trace elements (calcium, iron, fluoride, potassium, manganese and magnesium), tea polyphenols and catechins (especially EGCG = Epigallocatechingallate). The combination of these powerful ingredients provides an antioxidative (Chan et al., 1997), anti-inflammatory, cardiotonic (Hollman et al., 1997) and cancer preventive effects (Lambert et al., 2003). Green tea contributes to the dental health, reduces deposits in the blood vessels and protects against atherosclerosis (Chacko et

al., 2010). While the antioxidant properties are generally ascribed to the high content of tea polyphenols (Balentine et al., 1997), the positive influence on the body weight is attributable to the caffeine and catechins. The caffeine/catechins combination stimulates metabolism, increases energy expenditure, fat digestion and resorption in the stomach and the intestine (Bartista et al., 2009). Furthermore, the German Medical Journal reported in April 2010 that the catechins contained in green tea (EGCG) harmless "toxic" plaques in Alzheimer's.

Aloe Vera

Sumerian and Egyptian records occupied that the "plant of immortality" or the "blood of the gods" has been used medicinally 5,000 years ago. Scientists were able to detect more than 200 active ingredients till today. The Aloe Vera (Fig. 4) is naturally rich in polysaccharides, glycoproteins, vitamins, amino acids, enzymes and phytochemicals (such as essential oils, saponins, tannins, salicylic acid, sterols and aloin) in an unique pharmacological combination (Choi et al., 2001).



Fig. 4. Aloe Vera

Esua and Rauwald reported 2005 of a novel bioactive glucan, which attributed to the significant anti-inflammatory effect. The Aloe further harmonized the metabolism in all organs by regulation of the basic substances and the acid-base balance. In folk medicine the "wonderful desert lily" shows an excellent effect in inflammation (especially in the gastrointestinal tract), skin diseases (especially in Eczema, Dermatitis and Psorias), sunburns and wound raptures (Vogler, Ernst, 1999).

Edelweiss (*Leontopodii alpinum*) — Pure ingrediens, real effect! The most famous and most symbolic flower of the Alps is naturally rich in antioxidants, terpenes, caffee acid derivatives (Leontopodic acid) and Leoligin. So it is very popular in the cosmetic- (especially in UV-protective-and anti-aging products), food- and dietary supplement industries. In the Alpine region it is particularly known Edelweiss (Fig. 5) as "Bellyache flower" (such as abdominal pain, indigestion, diarrhea or intestinal colic)



Fig. 5. Edelweiss

The secondary plant substances of the "Edelweiss" support in the prevention and treatment of gastrointestinal complaints (Dobner et al., 2003) and in respiratory-, neurologic-, muscular- and cardiovascular diseases.

In addition to the outstanding antiinflammatory, antibacterial (Dobner et al. 2004) — even with infections (Stuppner, 2000) — and analgesic effects (Speroni et al., 2006) the "Edelweiss" shows a superior property as a scavenger and antioxidant (Schwalger et al., 2005) and preserves so cells from damage (also by molds; (Costa et al., 2009). While Leoligin (Fig. 6) proved highly effective against thickening of the inner wall of blood vessels (Reisinger et al., 2009), an unknown terpene shows a significant cancer hostile activity against human leukemia cells (Wang et al. 2007). Moreover (Hornick et al., 2008) can showed after taking an extract from the whole root an improvement in the capacity of memory and kick off a discussion for taking this extract in the prevention and treatment of dementia.

Fig. 6. The structure of Leoligin in Edelweiss

3.2. Comparative analysis between longevity of long living people, centenarians and their siblings and the quality of water

In frames of the research on the water quality 121 long living people from Bulgaria over 90 years of age have been studied together with their 294 siblings. The average lifespan of long lived people and centenarians in mountain areas is 94.1 years. For the average lifespan of long lived people in plain areas the result is 90.6 years. The most adult person from mountain areas is 104 years old and for plain areas is 97 years old. For the brothers and sisters of long live people from mountain areas the average lifespan is 88.5 years. For the brothers and sisters of long live people from plain areas the average lifespan is 86.4 years. The difference in life expectancy of the two groups of people is reliable and it corresponds to the Student's t-criteria at p < 0.05 with a confidence level of t = 2.36. There are distances of no more than 50-70 km between these places and the only difference is the mountain water and air.

There have been 21519 residents in Teteven and 142 of them were born before 1924. Figure 7 demonstrates the interrelation between the year of birth (1912–1924) of long living people (age) and their number (Teteven municipality, Bulgaria).




Fig. 7. Interrelation between the year of birth of long living people (age) and their number in Teteven municipality, Bulgaria

From the standpoint of genetics, the process of aging is associated with disruption of the genetic program of the organism and gradual accumulation of errors during the process of DNA replication. Aging may be associated with the accumulation of somatic mutations in the genome and be influenced by free radicals (mainly oxygen and primary products of oxidative metabolism) and ionizing radiation on DNA molecules as well (Woodhead, 1984; Adelman et al., 1988). Such mutations can reduce the ability of cells to the normal growth and division and be a cause of a large number of various cell responses: inhibition of replication and transcription, impaired cell cycle division, transcriptional mutagenesis, cell aging that finally result in cell death. Cells taken

from the elderly people show a reduction in transcription when transferring information from DNA to RNA.

From the standpoint of dynamics, aging is a non-linear biological process, which increases over time. Accordingly, the rate of aging increases with time. The accumulation of errors in the human genome increases exponentially with time and reaches a certain stationary maximum at the end of life. L. Orgel shows that, for this reason, the probability of cancer occurrence increases with age (Orgel, 1963). Figure 8 shows L. Orgel's results on the interrelation between age and the number of cancer cases. The accumulation of errors in synthesis of abnormal proteins increases exponentially over time with age. Cells taken from elderly people show the reduced levels of transcription or transmission of information from DNA to RNA. Therefore, the probability of cancer increases with age. The interrelation between the number of Bulgarian centenarians in the mountainous municipality of Teteven and their age is close to exponential.




Fig. 8. Interrelation between age and the number of cancer patients (Orgel, 1963)

Here are submitted the data for Bulgaria:

- 1) Varna district 44 centenarians per 1 million of inhabitants, plain and sea regions;
- 2) Pleven district 78 centenarians per 1 million of inhabitants, plain regions;
- 3) Teteven district 279 centenarians per 1 million of inhabitants, hills and mountainous regions;
 - 4) Bulgaria 47 centenarians per 1 million of inhabitants.

3.3. Clinical studies with human blood serum testing

A convenient method for studying of liquids is non-equilibrium differential spectrum. It was established experimentally that the process of evaporation of water drops, the wetting angle θ decreases discreetly to zero, and the diameter of the water drop basis is only slightly altered, that is a new physical effect (Antonov, 1995; Antonov, Yuskesselieva, 1983). Based on this effect, by means of the measurement of the wetting angle within equal intervals of time is determined the function of distribution of H₂O molecules according to the value of $f(\theta)$. The distribution function is denoted as the energy spectrum of the water state. The theoretical research established the dependence between the surface tension of water and the energy of hydrogen bonds among individual H₂O-molecules (Antonov, 1995).

For calculation of the function f(E) represented the energy spectrum of water, the experimental dependence between the wetting angle (θ) and the energy of hydrogen bonds (E) is established:

$$f(E) = \frac{14,33f(\theta)}{[1-(1+bE)^2]^2}$$
 (1)

where $b = 14.33 \text{ eV}^{-1}$

The relation between the wetting angle (θ) and the energy (E) of the hydrogen bonds between H₂O molecules is calculated by the formula:

$$\theta = \arccos\left(-1 - 14.33E\right) \tag{2}$$

The energy spectrum of water is characterized by a non-equilibrium process of water droplets evaporation, therefore, the term non-equilibrium spectrum (NES) of water is used.

The difference $\Delta f(E) = f(E \text{ samples of water}) - f(E \text{ control sample of water})$ – is called the "differential non-equilibrium energy spectrum of water" (DNES).

Thus, the DNES spectrum is an indicator of structural changes in water, because the energy of hydrogen bonds in water samples differ due to the different number of hydrogen bonds in water samples, which may result from the fact that different waters have different structures and composition and various intermolecular interactions – various associative elements etc. (Ignatov et al., 2014; Ignatov et al., 2015). The redistribution of H_2O molecules in water samples according to the energy is a statistical process of dynamics.

Figure 9 shows the average NES-spectrum of deionised water. On the X-axis are depicted three scales. The energies of hydrogen bonds among H_2O molecules are calculated in eV. On the Y-axis is depicted the function of distribution of H_2O molecules according to energies f(E), measured in reciprocal unit eV^{-1} .

Arrow A designates the energy of hydrogen bonds among H₂O molecules, which is accepted as most reliable in spectroscopy.

Arrow B designates the energy of hydrogen bonds among H₂O molecules the value of which is calculated as:

$$\bar{E} = -0.1067 \pm 0.0011 \text{ eV}$$
 (3)

Arrow C designates the energy at which the thermal radiation of the human body, considered like an absolute black body (ABB) with a temperature +36.6 °C, is at its maximum.

A horizontal arrow designates the window of transparency of the Earth atmosphere for the electromagnetic radiation in the middle infrared range of the Sun toward the Earth and from the Earth toward the surrounding space. It can be seen that the atmosphere window of transparency almost covers the NES-spectrum of water.




Fig. 9. The NES-spectrum of deionized water (chemical purity -99.99%; pH -6,5-7,5; total mineralization -200 mg/l; electric conductivity -10 μ S/cm): the horizontal axis shows the energy of the H...O hydrogen bonds in the associates -E (eV); the vertical axis - the energy distribution function -f (eV-1); k - the vibration frequency of the H-O-H atoms (cm-1); λ - wavelength (μ m)

We have conducted studies of 1 % (v/v) solution of human blood serum taken from two groups of people between 50 and 70 years of age by IR, NES and DNES spectral analysis. The first group consisted of people in excellent health. The second group consisted of people in a critical state and patients with malignant tumors. The average energy of hydrogen bonds ($\Delta E_{H...O}$) between H₂O molecules in the blood serum was investigated as the main biophysical parameter. The result was registered as a difference between the NES-spectrum of 1 % solution of human blood serum and the NES-spectrum of deionized water control sample - DNES-spectrum, measured as the difference $\Delta f(E) = f$ (samples of water) – f (control sample of water). The DNES-spectrum obtained from the first group has a local extremum energy ($\Delta E_{H...0}$) at E = -9.1±1.1 meV and from the second group at $E = -1.6 \pm 1.1$ meV. The results between the two groups have a statistical difference in Student's criterion at p < 0.05. For the control group of healthy people the value of the largest local maximum in the DNES-spectrum was detected at E = -0.1387 eV, or at a wavelength λ = 8.95 μ m. For the group of people in a critical health state and the patients with malignant tumors, the analogous values of the largest local maximums of the DNES-spectrum shifted to lower energies compared with the control group of people. For a group of people in critical health condition and patients with malignant tumors the greatest values of local extremum in the IR-spectrum are shifted to lower energies relative to the control group. In IR-spectrum of human blood serum are detected 8 local maxima at $\lambda = 8.55$, 8.58, 8.70, 8.77, 8.85, 9.10, 9.35 and 9.76 µm (Krasnov, Gordetsov, 2009). The resulting peak at $\lambda = 8.95 \mu m$ in the IR-spectrum (Ignatov, 2012) approaching the peak at $\lambda = 8.85 \mu m$ was monitored by Russian researchers. In the control group of healthy people the average value of the energy distribution function f(E) at $\lambda = 8.95 \mu m$ compiles E = 75.3 eV, and in a group of people in critical condition -E = 24.1 eV. The norm has statistically reliable result for human blood serum for the control group of people having cancer at the local extremum of f(E) ~24.1 eV-1. The level of reliability of the results is p< 0.05 according to the Student's t-test. In 1995 were performed DNES-experiments with an impact on tumor mice cells in water solutions containing Ca2+ (Antonov, 1995). There was a decrease in the DNES-spectrum compared with the control sample of cells from a healthy mouse. The decrease was also observed in the DNES-spectrum of human blood serum of terminally ill people relative to that of healthy people. With increasing of age of long-living blood relatives, the function of distribution of H₂O molecules according to energies at -0.1387 eV decreases. In this group of tested people the result was obtained by the DNES-method at $E = -5.5 \pm 1.1$ meV; the difference in age was of 20-25 years in relation to the control group. It should be noted that many of Bulgarian centenarians inhabit the Rhodopes Mountains areas. Among to the DNES-spectrum of mountain waters the similar to the DNES-spectrum of blood serum of healthy people at $\lambda = 8.95 \mu m$, was the DNES-spectrum of water in the Rhodopes. The mountain water from Teteven, Boyana and other Bulgarian provinces has similar parameters. Tables 1, 2 and 3 show the composition of mountain water springs in Teteven and Kuklen (Bulgaria) and local extremums in NES-spectra of water samples. The local extremums is water samples were detected at E = -0.11 eV and E = -0.1387 eV. The value measured at E = -0.11eV is characteristic for the presence of Ca²⁺ in water. The value measured at E = -0.1387 eV is characteristic for inhibiting the growth of cancer cells. Experiments conducted by A. Antonov with cancer cells of mice in water with Ca2+ demonstrated a reduction of this local extremum to a negative value in spectra. Analysis by the DNES-method of aqueous solutions of natural mineral sorbents – shungite (carbonaceous mineral from Zazhoginskoe deposit in Karelia, Russia) and zeolite (microporous crystalline aluminosilicate mineral from Most village, Bulgaria) showed the presence of a local extremum at E = -0.1387 eV for shungite and E = -0.11 eV for zeolite (Mosin, Ignatov, 2013, Ignatov, Mosin, 2014). It should be noted that owing to the unique porous structures both the natural minerals shungite and zeolite are ideal natural water adsorbers effectively removing from water organochlorine compounds, phenols, dioxins, heavy metals, radionuclides, and color, and gives the water a good organoleptic qualities, additionally saturating water with micro-and macro-elements until the physiological levels (Mosin & Ignatov, 2013). It is worth to note that in Bulgaria the main mineral deposits of Bulgarian zeolites are located in the Rhodope Mountains, whereat has lived the greatest number of Bulgarian centenarians. It is believed that water in these areas is cleared out in a natural way by mineral zeolite. Company LavaVitae has medicine product with high quality ZEOLITH detox.

3.4. Results of 1 % (v/v) solution in deionized water of VITA intense

The research with the NES method of water drops is received with 1 % solution VITA intense, and deionized water as control sample. The mathematical models of 1 % (v/v) solution VITA intense gives the valuable information for the possible number of hydrogen bonds as percent of H_2O molecules with different values of distribution of energies (Table 2 and Fig. 10). These distributions are basically connected with the restructuring of H_2O molecules having the same energies.

Table 2. The distribution (%, (-E_{value})/(-E_{total value}) of H₂O molecules in 1 % water solution of VITA intense (product of LavaVitae, Austria) and control deionized water

| -E(eV) | 1 % water | Control | -E(eV) | 1 % water | Control |
|--------|--------------------------------|----------------------------------|--------|--------------------------------|--------------------------------|
| x-axis | solution | Sample | x-axis | solution | Sample |
| | VITA intense | Deionized | | VITA intense | Deionized water |
| | (LavaVitae) | water | | (LavaVitae) | y-axis |
| | y-axis | y-axis | | y-axis | $(\%((-E_{value})^*/$ |
| | (%((-E _{value}) */ | $(\%((-E_{value})^*/$ | | $(\%((-E_{value}) */$ | $(-E_{\text{total value}})$ ** |
| | (-E _{total value})** | $(-E_{\text{total value}})^{**}$ | | (-E _{total value})** | |
| 0.0937 | 0 | 0 | 0.1187 | 0 | 0 |
| 0.0962 | 0 | 14.3 | 0.1212 | 17.7^{2} | 0 |
| 0.0987 | 0 | 0 | 0.1237 | 0 | 0 |
| 0.1012 | 0 | 14.3 | 0.1262 | 11.7 | 3.6 |
| 0.1037 | 0 | 0 | 0.1287 | 0 | 3.6 |
| 0.1062 | 11.7 | 14.3 | 0.1312 | 5.7 | 7.1 |
| 0.1087 | 0 | 0 | 0.1337 | 11.7 | 0 |
| 0.1112 | 5. 7¹ | 14.3 | 0.1362 | 5.7 | 7.1 |
| 0.1137 | 0 | 0 | 0.1387 | 30.1 ³ | 7.1 |
| 0.1162 | 0 | 14.3 | _ | _ | _ |

E= -0.1112 eV is the local extremum for relaxing effect on nervous system

E=-0.1212 eV is the local extremum for anti-inflammatory effect

E= -0.1387 eV is the local extremum for inhibition of development of tumor cells of molecular level

Notes:

* The result (-E_{value}) is the result of hydrogen bonds energy for one parameter of (-E)

** The result (-E_{total value}) is the total result of hydrogen bonds energy




Fig. 10. Mathematical model (Ignatov, Mosin, 2013) of 1 % water solution of VITA intense (product of LavaVitae, Austria).

Figure 10 shows the distribution (%, $(-E_{value})/(-E_{total\ value})$ of H_2O molecules in and 1 % (v/v) of water solution of VITA intense (product of LavaVitae, Austria) (red line) and control sample deionized water (blue line).

Notes:

E= -0.1112 eV is the local extremum for relaxing effect on nervous system

E=-0.1212 eV is the local extremum for anti-inflammatory effect

E= -0.1387 eV is the local extremum for inhibition of development of tumor cells of molecular level

The experimental data obtained testified the following conclusions from the mathematical model of in 1 % (v/v) water solution of VITA intense (product of LavaVitae, Austria) and control deionized water. The distribution (%, (-E_{value})/(-E_{total value}) of water molecules in mathematical model of in 1 % (v/v) water solution of VITA intense (product of LavaVitae, Austria) and control deionized water. The distribution (%, (-E_{value})/(-E_{total value}) of water molecules in VITA intense (product of Lava Vitae, Austria) according control sample is different. However, for the value E = -0.1387 eV or $\lambda = 8.95 \, \mu m$ there is the biggest local extremum (30.1 (%, (-E_{value})/(-E_{total value})) corresponding to the re-structuring of hydrogen bonds among H₂O molecules for inhabitation of development of tumor cells of molecular level. This difference may indicate on the different number of hydrogen bonds in water samples, as well as their physical parameters (pH, ORP), resulting in different distribution of H₂O molecules and different values of H₂O molecules with ratios of (-E_{value})/(-E_{total value}). Particularly it was observed the statistical re-structuring of H₂O molecules in water samples according to the energies. The experimental data may prove that stipulates the restructuring of H₂O molecules on molecular level and may be used for the prophylaxis of inhibition of development of tumor cells. For the value E = -0.1112 eV or $\lambda = 11.15$ um there is the local extremum with sight minus (5.7 (%, (-E_{value})/(-E_{total value})) according control sample (14.3 (%, (-E_{value})/(-E_{total value})) corresponding to the re-structuring of hydrogen bonds among H₂O molecules. The experimental data may prove that influence stipulates the restructuring of H₂O molecules on molecular level and has biophysical effect for relaxing effect on nervous system. For the value E = -0.1212 eV or λ = 10.23 μ m there is bigger local extremum (17.7 (%, (-E_{value})/(-E_{total value})) corresponding to the re-structuring of hydrogen bonds among H₂O molecules for anti-inflammatory effect. The experimental data for Lava Intense may prove that stipulates the restructuring of H₂O molecules on molecular level and the biophysical effects are:

E= -0.1112 eV is the local extremum for relaxing effect on nervous system

E=-0.1212 eV is the local extremum for anti inflammatory effect

E= -0.1387 eV is the local extremum for inhibition of development of tumor cells of molecular level.

As a result of different energies of hydrogen bonds, the surface tension of 1% (v/v) solutions of water samples with VITA Intense is increasing. The increasing of surface tension is regarding the control samples. This effect is connected with preservation of the energy in human body as result of biochemical process among water molecules and bio molecules.

3.5. VITA intense (product of the company LavaVitae)

The average energy (E_{H...O}) of hydrogen H...O-bonds among individual H₂O molecules in 1 % (v/v) solution of VITA Intense is measured at E=-0.1261 eV. The result for the control sample (deionized water) is E=-0.1125 eV. The results obtained with the NES method are recalculated with the DNES method as a difference of the NES (1 % (v/v) solution of VITA intense) minus the NES (control sample with deionized water) equaled the DNES spectrum of 1 % solution of VITA Intense. Thus, the result for 1 % solution of VITA intense recalculated with the DNES method is $\Delta E=$ 0.0136±0.0011 eV. The result shows the increasing of the values of the energy of hydrogen bonds in 1 % (v/v) solution of VITA intense regarding the deionized water. This is effect of stimulation on human body. The results show restructuring of water molecules in configurations of clusters, which influence usefully on human health on molecular and cellular level. The effects are describing with mathematical model of 1 % solution of VITA intense.

 ${\it 3.6. Results with pH and ORP}$ There are valid the following results of pH as indicator for acid alkaline medium of the products of LavaVitae. There are the results also of ORP or Oxidation-reduction potential.

The results are for 1% (v/v) of solutions of products, which are made from deionized water. This research is performed with Georgi Gluhchev from Bulgarian Academy of Science. The results of pH of deionized water is 6.05 and of ORP is 119.7. Table 3 shows the results of pH and ORP.

| Table 3. Results | of products | of company Lava | Vitae for pH and ORP |
|-------------------------|-------------|-----------------|----------------------|
|-------------------------|-------------|-----------------|----------------------|

| Product | рН | ORP (mV) | Coordinates Fig. 11 |
|--------------------|-----------|----------|---------------------------|
| VITA intense | 4.07±0.02 | - 104.5 | Point 1 (4,07; -104.5) |
| BOOST | 3.60±0.02 | +113.6 | Point 2 (3,90;113.6) |
| ZEOLITH detox | 8.01±0.02 | +109.5 | Point 3 (8,01;103.3) |
| Deionized water | 6.05±0.02 | +119.7 | |

Figure 11 shows the dependence between the acidity and basicity (pH) of electrochemically activated solutions and the oxidation-reduction potential (ORP). The pH value within the interval from 3 to 10 units and the ORP within the interval from -400 mV to +900 mV characterize the area of the biosphere of microorganisms. Outside these ranges of pH and ORP the microorganisms will hardly survive.




Fig. 11. The dependence between acidity and basicity (pH) of solutions and the ORP on the biosphere of micro-organisms (point 1; VITA intense), (point 2; BOOST), point 3; ZEOLITH detox)

The result of 1 % (v/v) solution of VITA intense is 4.07 or acidic medium. The result of ORP is (-104.5). The result of ORP with negative charge is connected with charge with negative value, which has antioxidant and permanent antioxidant activity. In the VITA intense there are the following antioxidants – Vitamins C, E, D. Figure 11 shows the dependence between acidity and basicity (pH) of solutions and the ORP on the biosphere of micro-organisms. The result of VITA intense with point 1 with coordinates (4,07; -104.5) is the biosphere of micro-organisms. VITA intense is useful for human health also with liquid form.

3.7. Effects of Magnesium and Selenium, Vitamins C and E in VITA intense again oxidative stress

VITA intense is product with negative Oxidation-reduction potential (ORP) or (-104.5) mV. This results shows that there are particles in VITA intense with negative charge again free radicals.

There are also active antioxidants as Vitamin C and Vitamin E. There are special role in the liquid product the ions of Magnesium – Mg²⁺. These ions neutralize one of the most danger radicals – hydroxyl radical OH (Glamello et al., 1993). Magnesium deficiency increased cytotoxicity of the added oxyradicals. These results suggest that increased oxidative endothelial cell injury may contribute to vascular injury during Mg deficiency (Dickens et al., 1992). The magnesium deficiency and oxidative stress have both been identified as pathogenic factors in aging and in several age-related diseases. The link between these two factors is unclear in humans although, in experimental animals, severe Mg²⁺ deficiency has been shown to lead to the increased oxidative stress (Begona et al., 2000). A negative correlation between magnesium balance and oxidative stress was observed suggesting that the same etiological factor (chronic stress) initiate decreases in both free and total magnesium concentrations and simultaneously increase oxidative stress intensity. These findings support the need for magnesium supplementation with antioxidant vitamins for people living in conditions of chronic stress (Cernak et al., 2000).

Selenium has role for anti oxidative stress in the enzymes, which are connected with anti oxidant effects (Stewart et al., 1999).

Vitamin E, as a powerful antioxidant residing mainly in biomembranes, may provide effective protection against oxidative membrane damage and resultant age-related deterioration, especially in the elderly. We hypothesized that appropriate levels of vitamin E supplementation would protect erythrocyte membranes from oxidative stress and thus improve membrane fluidity in healthy middle-aged and elderly people (Sun et al., 2012).

The combination of antioxidative treatment with vitamins E and C decreases fetal malformation rate and diminishes oxygen radical-related tissue damage (Cederberget al., 2001).

3.8. Effects of Ca^{2+} , Mg^{2+} , Zn^{2+} and Mn^{2+} in water on biophysical and biochemical processes in the human body

The research of distribution of local extremums (eV-1) in spectra of various water samples as a function of distribution of H₂O molecules according to energy f(E) at $\lambda = 8.95 \mu m$ shows the analogue extremum at analogous values of f(E), E and λ, which was detected in water with Zn²⁺and Mn²⁺ ions earlier demonstrated inhibiting the growth of cancer cells. Magnesium (Mg²⁺), zinc (Zn²⁺) and manganese (Mn²⁺) ions dissolved in water have influence on enzymes, which are antioxidants (Ignatov, Mosin, 2015). The research of China team was categorized three groups of elements from the rice and drinking water according to their effect on longevity: Sr, Ca, Al, Mo, and Se, which were positively correlated with longevity: Fe, Mn, Zn, Cr, P, Mg, and K, which had a weak effect on local longevity, and Cu and Ba, which had a negative effect on longevity (Lv et al., 2011). There was a positive correlation between the eSOD activity and the age and a negative correlation between the eSOD activity and concentration of Zn²⁺ in plasma. An inverse correlation was also found between the content of Zn²⁺ ions in plasma relative to the age. The prevalence of Zn²⁺ deficiency is increased with age; with normal Zn²⁺ levels it is observed in about 80 % of adult people and only in 37 % of the non-agenarians. Aging is an inevitable biological process that is associated with gradual and spontaneous biochemical and physiological changes and the increased susceptibility to diseases. Because the nutritional factors are involved in improving the immune functions, metabolic balance, and antioxidant defense, some nutritional factors, such as Zn, may modify susceptibility to disease and promote healthy aging. In vitro (human lymphocytes exposed to endotoxins) and in vivo (old or young mice fed with low zinc dietary intake) studies revealed that zinc is important for immune efficiency (innate and adaptive), antioxidant activity (superoxide dismutase), and cell differentiation via clusterin/apolipoprotein J. The intracellular Zn homeostasis is regulated by metallothioneins (MT) via an ion release through the reduction of thiol groups in the MT molecule (Mocchegiani, 2007). Zinc in composition of water improves the antioxidative enzymes in red blood cells (Malhotra, Dhawan, 2008).

The antioxidants against free radical damage include tocopherol (vitamin E), ascorbic acid (vitamin C), β -carotene, glutathione, uric acid, bilirubin, and several metalloenzymes including glutathione peroxidase (Se), catalase (Fe), and superoxide dismutase (Cu, Zn, Mn) and proteins such as ceruloplasmin (Co). The extent of the tissue damage is the result of the balance between the free radicals generated and the antioxidant protective defense system (Machlin, Bendich, 1988). There was reported the antioxidant effects of water on rats (Abdullah, 2012). The norm in water for

 Zn^{2+} and Mn^{2+} according to the World Health Organization (WHO) should be less than 20 μ g. For the Na⁺ content the norm according to the WHO is less than 20 μ g.

The interesting results on the concentration of Ca²⁺ in water were obtained in USA and Canada. According to the statistical information the most number of centenarians in Canada per 1 million of population is observed in Nova Scotia (210 of centenarians per 1 million). In the water from Nova Scotia the Ca²⁺ content makes up 6.8 mg/l. N. Druzhyak, Russia showed that in the places wherein live the most number of centenarians the Ca²⁺ content in water was 8–20 mg/l. The only risk factor regarding the increased Ca²⁺ content in water is cardiovascular diseases.

The following reactions occur in water if there are high concentrations of Ca^{2+} and Mg^{2+} ions: the reaction of limestone ($CaCO_3$) and gypsum ($CaSO_4 \cdot 2H_2O$) with water to separate the calcium (Ca^{2+}), carbonates (CO_3^{2-}) and sulfate (SO_4^{2-}) ions. By increasing the mineralization of water the content of Ca^{2+} ions decreases. During the concentration of the solutions Ca^{2+} ions are precipitated. With the increase of carbon dioxide (CO_2) in water and decreasing of the pH value the content of Ca^{2+} increases. The reaction of interaction of dolomite ($CaCO_3 \cdot MgCO_3$) with water makes the formation of Ca^{2+} ions. Hydrocarbonates ($CO_3 \cdot MgCO_3$) ions are formed by reaction of interaction of karst rocks, CO_2 and water. For example, in Zamzam water there is $Ca^{2+} - 299.7 \cdot mg/l$; $CaCO_3 \cdot MgCO_3 \cdot MgCO_3$

4. Conclusion

From the NES and DNES spectrum and mathematical model of 1% (v/v) solution of VITA Intense and deionized water as control sample are valid the following conclusions for biophysical effects for VITA Intense (LavaVitae company)

- relaxing effect on nervous system;
- anti inflammatory effect;
- inhibtion of development of tumor cells of molecular level;

In 1 % (v/v) solution of VITA intense there is restructuring of water molecules in configurations of clusters, which influence usefully on human health on molecular and cellular level.

The biophysical effects of VITA intense are connected also with antioxidant effects. The VITA intense is recommended as anti aging solution for prophylaxis (Ignatov et al., 2015). The scientific studies show that the inflammations are one of the basic reasons for aging. The recommendation is connected with additionally using or including in VITA intense of the additional mineral. The structuring of water clusters with highest energy of hydrogen bonds at 8.95 µm makes the water in human body more "active" as medium of biochemical and biophysical processes. This is similar like the human organism to be younger (Ignatov, Mosin, 2012). The quality of the water with which will be using is very important. There are types of water which will increase the effects. For these effects is recommended additional scientific project with pH is in progress.

As a result of different energies of hydrogen bonds, the surface tension of 1% (v/v) solutions of water samples with VITA intense is increasing. The increasing of surface tension is regarding the control sample. This effect is connected with preservation of the energy in human body as result of biochemical process among water molecules and bio molecules;

It worth to note that IR-spectrum of VITA intense is most similar to the IR-spectrum of blood serum of healthy group of people with a local extremum at $\lambda=8.95$ µm. The similar spectral characteristics possess mountain water from Teteven and other Bulgarian sources electrochemically activated water catholyte (Ignatov et al., 2014). The basic research is with electrolysis device (Ignatov et al., 2014) Studying the human blood serum by NES and DNESmethods show that by measuring the average energy of hydrogen bonds among H₂O molecules and the distribution function of H₂O molecules on energies it is possible to show a vital status of a person and associated life expectancy. Our data indicates that water in the human body has the IRspectrum resembling the IR-spectrum of human blood serum. On the characteristics of the IRspectrum of water also exerts an influence the presence of deuterium in water samples. In the research there is the optimal composition of mountain and melt water from areas where are lived the long live people and centenarians. The decreased content of deuterium in studied water samples with residual deuterium content of 60-100 ppm, the variety of ions (K⁺, Na⁺, Ca²⁺, Mg²⁺, Mn²⁺, Fe²⁺, Fe³⁺, Zn²⁺, SO₄²⁻, Cl⁻, HCO₃⁻, CO₃²⁻), and chemical-physical parameters (pH, electroconductivity) of studied water samples renders beneficial effects of this type of water on human health. We have also obtained new proofs for biophysical and biochemical effects of Ca²⁺, Mg²⁺, Zn²⁺ and Mn²⁺ in composition of water on human organism and DNES-spectra of water. There are obtained new results of chemical composition of water from Glacier Rosenlaui, Swiss Alps.

VITA intense is an elixir from alpine herbs that strengthens the body powerfully. It is most useful during the autumn, winter and spring seasons, when it is the time of the viruses. It contains a complex of edelweiss, aronia, aloe vera and green tea. And most importantly is, that the antioxidant vitamins C and E, as well as Magnesium are together in a liquid that gets absorbed from the body perfectly. The magnesium neutralizes one of the most dangerous radicals, explains the scientist – the hydroxyl radical OH. This radical is one of the reason for Parkinson, Alzheimer and whole status of the brain and nervous system. VITA intense is one of the not so many liquid products from this type, which imports into the body negative antioxidant charges with effects again tumor cells. Inflammations are basic reason for aging process in human body. In VITA intense there are the following effects – anti aging, relaxing effect on the nervous system, antioxidant, inhibition of development of tumor cells, stimulating on the body. The people take one product with a lot of effects. VITA intense id in glass bottle and this is very important for the quality. People feel rush of energy, life vigour and a sense of happiness.

References

Abdullah et al., 2012 – Abdullah, A.M., Abdelsalam, E., Abdullah, B., Khaled, A. (2012). Antioxidant Effects of Zamzam Water in Normal Rats and Those Under Induced-oxidant Stress, *Journal of Medicinal Plants Research*, 6(42), 5507-5512.

Adelman et al., 1988 – Adelman, R., Saul, R., Ames, B. (1988). Oxidative Damage to DNA: Relation to Species Metabolic Rate and Life Span, PNAS, 85(8), 2706-2708.

Antonov, 1995 – Antonov, A. (1995). Research of the Non-equilibrium Processes in the Area in Allocated Systems, *Dissertation Thesis for Degree "Doctor of Physical Sciences"*, Blagoevgrad, Sofia.

Antonov, Yuskesselieva, 1985 – Antonov, A., Yuskesselieva, L. (1985). Selective High Frequency Discharge (Kirlian Effect), Acta Hydrophysica, 5, 29.

Atanasov et al., 2014 – Atanasov, A., Karadzhov, S., Ivanova, E., Mosin, O.V., Ignatov I. (2014). Study of the Effects of Electrochemical Aqueous Sodium Chloride Solution (Anolite) on the Virus of Classical Swine Fever Virus. Mathematical Models of Anolyte and Catolyte as Types of Water, Journal of Medicine, Physiology and Biophysics, 4, 1–26.

Begona et al., 2000 – Begona, M. et al. (2000). Magnesium Status and Parameters of the Oxidant-Antioxidant Balance in Patients with Chronic Fatigue: Effects of Supplementation with Magnesium, Journal of American College for Nutrition, 19 (3), 374-382.

Balentine et al., 1997 – Balentine, D. et al. (1997). The Chemistry of Tea Flavonoids, Food Science and Nutrition, 37 (8), 693-70.

Bell, Goshenaur, 2006 – *Bell, D.R., Goshenaur, K.* (2006). Direct Vasodilatative Properties of Anthocyanin-Rich Extract, *J. Appl. Physiol.*, 100, 1164-1170.

Batista et al., 2009 – *Batista et al.* (2009). Prospective Double-blind Crossover Study of Camellia Sinensis (Green tea) in Dyslipidemias, *Arq. Bras. Cardiol.*, 93 (2), 128-134.

Bermudez-Soto et al., 2007 – Bermudez-Soto et al. (2007). Stability of Polyphenols in Chokeberry (Aronia melanocarpa) Subjected to in vitro Gastric and Pancreatic Digestion, Food Chemistry, 102 (3), 865-874.

Borisova et al., 2004 – *Borisova et al.* (2004). Hepatoprotective Effect of the Natural Fruit Juice from Aronia melanocarpa on Carbon Tetrachloride-induced Acute Liver Damage in Rats, *Exp. Toxicol. Pathol.*, 56 (3), 195-201.

Brubach et al., 2005 – Brubach, J.B., Mermet, A., Filabozzi, A., Gerschel, A., Roy, P. (2005) Signatures of the Hydrogen Bonding in the Infrared Bands of Water, J. Chem. Phys., 122.

Burger et al., 2012 – Burger, O., Baudish, A. &Vaupel, J. W. (2012). Human Mortality Improvement in Evolutionary Context, PNAS, 109(44), 18210–18214.

Chan E-V et al., 2011 – Chan E-V et al. (2011). Antioxidant and Antibacterial Properties of Green, Black, and Herbal Teas of Camellia sinensis, Pharmacognocy Research, 3(4), 266-272.

Chacko et al., 2010 – Chacko, S. et al. (2010). Beneficial Effects of Green Tea. A literature Review, Chinless Medicine, 5-13.

Choi et al., 2001 – Choi et al. (2001). The Green Tea Polyphenol(-)-epigallocatechin Gallate Attenuates Beta-amyloid-induced Neurotoxicity in Cultured Hippocampal Neurons, *Life Sci.*, 70 (5), 603-614.

Chan, 1997 – Chan M-Y (1997). Inhibition of Inducible Nitric Oxide Synthase Gene Expression and Enzyme Activity by Epigallocatechin Gallate, a Natural Product from Green Tea, *Biochemical Pharmacology*, 54 (12), 1281–1286.

Cederberg et al., 2001 – Cederberg, J., Siman, C., Eriksson, U. (2001). Combined Treatment with Vitamin E and Vitamin C Decreases Oxidative Stress and Improves Fetal Outcome in Experimental Diabetic Pregnancy, *Pediatric Research*, 49, 755–762.

Costa et. al., 2009 – Costa, S. et. al. (2009). In vitro Evaluation of the Chemoprotective Action Mechanisms of Leontopodic Acid Against Aflatoxin B1 and Deoxynivalenol-induced Cell Damage, J Appl Toxicol., 29 (1),7-14.

Choi, Jordan, 2010 – Choi T.N., Jordan K.D. (2010). Application of the SCC-DFTB Method to $H^+(H_2O)_{6}$, $H^+(H_2O)_{21}$, and $H^+(H_2O)_{22}$, J. Phys. Chem. B, 114, 6932–6936.

Cernak et al., 2000 – Cernak et al. (2000). Alterations in Magnesium and Oxidative Status during Chronic Emotional Stress, Magnesium Research, 13(1), 29-36.

Davis, 1989 – Davis, R.H. (1989). Wound Healing, Oral & Topical Activity of Aloe Vera, J. Am. Podiatr. Med. Assoc., 79 (11), 559-562.

Dickens et al., 1992 – *Dickens, B. et al.* (1992) Magnesium Deficiency in vitro Enhances Free Radical-induced Intracellular Oxidation and Cytotoxicity in Endothelial Cells, *FEBS letters*, 13 (2), 102–108.

Dobner et al., 2003 – Dobner, M. et al. (2003). Antibacterial Activity of Leontopodium alpinum (Edelweiss), Journal of Ethnopharmacology, 89 (2-3), 301-303.

Druzhyak, 2005 – Druzhyak, N.G. (2005). Water for Health and Longevity, Neva. [in Russian]

Eisenberg, Kauzmann, 1969 – Eisenberg, D., Kauzmann, W. (1969). The Structure and Properties of Water, Oxford University Press, London.

Esua, Rauwald, 2006 – Esua, M., Rauwald, J-W. (2006). Novel Bioactive Maloyl Glucans from Aloe vera Gel: Isolation, Structure Elucidation and in Vitro Bioassays, Carbohydrate Research, 341 (3), 355-364.

Glamello et al., 1993 – Glamello, E. et al. (1993). Evidence of Stable Hydroxyl Radicals and Other Oxygen Radical Species Generated by Interaction of Hydrogen Peroxide with Magnesium Oxide, *The Journal of Physical Chemistry*, 97 (21), 5735–5740.

Gluhchev et al., 2015a — Gluhchev, G., Ignatov, I. Karadzhov, S., Miloshev, G., Ivanov, N., Mosin, O.V. (2015). Electrochemically Activited Water. Biophysical and Biological Effects of Anolyte and Catholyte as Types of Water, Journal of Medicine, Physiology and Biophysics, 10, 1-17.

Gluhchev et al., 2015b – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, N., Mosin, O.V. (2015). Studying the Antimicrobial and Antiviral Effects of Electrochemically Activated Nacl Solutions of Anolyte and Catholyte on a Strain of E. Coli DH5 and Classical Swine Fever (CSF) Virus, European Journal of Medicine, 9 (3), 124-138.

Gluhchev et al., 2015c – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O.V. (2015). Studying of Virucidal and Biocidal Effects of Electrochemically Activated Anolyte and Catholyte Types of Water on Classical Swine Fever Virus (CSF) and Bacterium E. coli DH5, Journal of Medicine, Physiology and Biophysics, 13: 1-17.

Gluhchev et al., 2015d – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O.V. (2015). Biocidal Effects of Electrochemically Activated Water, Journal of Health, Medicine and Nursing, 11, 67-83.

Gluhchev et al., 2015e – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, N., Mosin, O.V. (2015). Antimicrobial and Antiviral Effects of Electrolytic Water Sodium Chloride (Anolyte/Catholyte) on a Strain of E.Coli DH5 and the Classical Swine Fever Virus (CSF), Questions of Medical, Biological and Pharmaceutical Chemistry, 3, 21-31.

Gluhchev et al., 2015f – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, N., Mosin, O.V. (2015). Electrochemically Activited Water: Biophysical and Biological Effects of Anolyte and Catholyte Types of Water, European Journal of Molecular Biotechnology, 7 (1), 12-26.

Gluhchev et al., 2015g – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, N., Mosin, O.V. (2015). Virucidal and Bactericidal Effect of Anolyte and Catholyte Types of Water of

Classical Swine Fever Virus and Bacteria E. Coli DH5, Water: Hygiene and Ecology, 3 (1-2): 5-23.

Goryainov, 2012 – Goryainov S.V. (2012). A Model of Phase Transitions in Double-well Morse Potential, *Application to Hydrogen Bond. Physica B*, **407**, 4233–4237.

Hollman et al., 1997 – *Hollman, P., Tjiburg, L., Yang, C.* (1997). Bioavailability of Flavonoids from Tea, *Crit. Rev. Food Sci. Nutr.*, 37, 719-738.

Howard, Hopps, 1986 – Howard, C., Hopps (1986). Chemical Qualities of Water that Contribute to Human Health in a Positive Way, *Science of the Total Environment*, 54, 207-216.

Ignatov et al., 2015 – Ignatov, I., Mosin, O.V., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, N. (2015). The Evaluation of Mathematical Model of Interaction of Electrochemically Activated Water Solutions (Anolyte and Catholyte) with Water, European Reviews of Chemical Research, 2 (4), 72-86.

Ignatov, 2005 – Ignatov, I. (2005). Energy Biomedicine, Gea-Libris, Sofia, 1–88.

Ignatov, 2010 – *Ignatov*, *I*. (2010). Which Water is Optimal for the Origin (Generation) of Life? *Euromedica*, Hanover, 34-35.

Ignatov, 2011 – Ignatov, I. (2011). Entropy and Time in Living Matter, Euromedica, 74.

Ignatov, 2012 – *Ignatov, I.* (2012). Origin of Life and Living Matter in Hot Mineral Water, Conference on the Physics, Chemistry and Biology of Water, *Vermont Photonics*, USA.

Ignatov, Mosin, 2013a – Ignatov I., Mosin O.V. (2013). Possible Processes for Origin of Life and Living Matter with Modeling of Physiological Processes of Bacterium *Bacillus Subtilis* in Heavy Water as Model System, *Journal of Natural Sciences Research*, 3 (9), 65-76.

Ignatov, Mosin, 2013b – Ignatov, I., Mosin, O. V. (2013). Modeling of Possible Processes for Origin of Life and Living Matter in Hot Mineral and Seawater with Deuterium, Journal of Environment and Earth Science, 3(14), 103-118.

Ignatov, Mosin, 2013c – *Ignatov, I., Mosin, O.V.* (2013). Structural Mathematical Models Describing Water Clusters, *Journal of Mathematical Theory and Modeling*, 3 (11), 72-87.

Ignatov, Mosin, 2014a – *Ignatov, I., Mosin, O. V.* (2014). The Structure and Composition of Carbonaceous Fullerene Containing Mineral Shungite and Microporous Crystalline Aluminosilicate Mineral Zeolite. Mathematical Model of Interaction of Shungite and Zeolite with Water Molecules., *Advances in Physics Theories and Applications*, 28, 10-21.

Ignatov, Mosin, 2014b – *Ignatov, I., Mosin, O.V.* (2014). The Structure and Composition of Shungite and Zeolite. Mathematical Model of Distribution of Hydrogen Bonds of Water Molecules in Solution of Shungite and Zeolite. *Journal of Medicine, Physiology and Biophysics*, 2, 20-36.

Ignatov, Mosin, 2014c – *Ignatov*, *I., Mosin,O.V.* (2014). Mathematical Models of Distribution of Water Molecules Regarding Energies of Hydrogen Bonds, *Journal of Medicine*, *Physiology and Biophysics*, 2, 71-94.

Ignatov, Mosin, 2014d – *Ignatov*, *I., Mosin,O.V.* (2014). Mathematical Model of Interaction of Carbonaceous Fullerene Containing Mineral Shungite and Aluminosilicate Mineral Zeolite with Water, *Journal of Medicine*, *Physiology and Biophysics*, 3, 15-29.

Ignatov, Mosin, 2014e – Ignatov, I., Mosin, O. V., Bauer, E. (2014). Carbonaceous Fullerene Mineral Shungite and Aluminosilicate Mineral Zeolite. Mathematical Model ans Practical Application of Water Solution of Water Shungite and Zeolite, Journal of Medicine, Physiology and Biophysics, 4, 27-44.

Ignatov, Mosin, 2016a – *Ignatov, I., Mosin, O.V.* (2016). Water for Origin of Life, *Altaspera Publishing & Literary Agency Inc*, 1-616. [in Russian]

Ignatov, Mosin, 2016b – *Ignatov, I., Mosin, O.V.* (2016). Deuterium, Heavy Water and Origin of Life, *LAP LAMBERT Academic Publishing*, 1-500.

Ignatov et al., 2016a – *Ignatov, I. et al.* (2016). Results of Biophysical and Nano Technological Research of ZEOLITH Detox of LavaVitae Company, *Journal of Health, Medicine and Nursing*, 30, 44-49.

Ignatov et al., 2016b – *Ignatov*, *I.* (2016). Product of LavaVitae BOOST is Increasing of Energy of Hydrogen Bonds among Water Molecules in Human Body, *Journal of Medicine*, *Physiology and Biophysics*, 27., 30-42.

Ignatov, 2016 – *Ignatov*, *I*. (2016). VITA intense – Proofs for Anti-inflammatory, Antioxidant and Inhibition Growth of Tumor Cells Effects. Relaxing Effect of Nervous System, Anti Aging Influence, *Journal of Medicine, Physiology and Biophysics*, 27, 43-61.

Ignatov, 2017a - Ignatov, I. (2017). VITA intense and BOOST - Products with Natural

Vitamins and Minerals for Health, Journal of Medicine, Physiology and Biophysics, 31, 58-78.

Ignatov, 2017b – Ignatov, I. (2017). ZEOLITH detox for Detoxification and ZELOLITH Creme for Skin Effects as Products of LavaVitae Company, Journal of Medicine, Physiology and Biophysics, 31, 79-86.

Ignatov, 2017c – *Ignatov*, *I*. (2017). Distribution of Molecules of ZEOLITH detox and ZEOLITH Creme in Water as Factor for Health, *European Journal of Molecular Biotechnology*, 5, (1), 11-22.

Ignatov, 2017d – *Ignatov*, *I.* (2017). Biophysical Research of ZEOLITH detox and ZEOLITH Creme, *European Journal of Medicine*, 5 (2), 31-42.

Ignatov, 2017e – *Ignatov*, *I*. (2017). VITA intense –Antioxidant and Inhibition Growth of Tumor Cells Effects. Anti Aging Influence. Negative Oxidation-reduction Potential (ORP) Has Important Role in These Effects, *Journal of Medicine*, *Physiology and Biophysics*, 39, 20-42.

Ignatov et al., 2014 – Ignatov, I., Karadzhov, S., Atanasov, A., Ivanova, E., Mosin, O.V. (2014). Electrochemical Aqueous Sodium Chloride Solution (Anolyte and Catholyte) as Types of Water. Mathematical Models. Study of Effects of Anolyte on the Virus of Classical Swine Fever Virus, Journal of Health, Medicine and Nursing, 8, 1-28.

Ignatov et al., 2015a – Ignatov, I., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O.V. (2015). Preparation of Electrochemically Activated Water Solutions (Catholyte/Anolyte) and Studying Their Physical-Chemical Properties, Journal of Medicine, Physiology and Biophysics, 11, pp. 1-21.

Ignatov et al., 2015b – *Ignatov, I., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O.V.* (2015). Preparation of Electrochemically Activated Water Solutions (Catholyte/Anolyte) and Studying of their Physical-Chemical Properties, *Journal of Medicine, Physiology and Biophysics*, 13, 18-38.

Ignatov et al., 2015c – Ignatov, I., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O.V. (2015). Preparation of Electrochemically Activated Water Solutions (Catholyte/Anolyte) and Studying of their Physical-Chemical Properties, Journal of Health, Medicine and Nursing, 13, 64-78.

Ignatov et al., 2015d – Ignatov, I., Mosin, O. V., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I. (2015). Studying Electrochemically Activated NaCl Solutions of Anolyte and Catholyte by Methods of Non-Equilibrium Energy Spectrum (NES) and Differential Non-Equilibrium Energy Spectrum (DNES), Journal of Medicine, Physiology and Biophysics, 14, 6-18.

Ignatov et al., 2015e – Ignatov, I, Gluhchev, G., Karadzhov, S., Ivanov, N., Mosin, O.V. (2015). Preparation of Electrochemically Activated Water Solutions (Catholyte/Anolyte) and Studying Their Physical-Chemical Properties, Journal of Medicine, Physiology and Biophysics, 16, 1-14.

Ignatov et al., 2016 – Ignatov, I., Mosin, O.V., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I. (2016). Studying Electrochemically Activated NaCl Solutions of Anolyte and Catholyte by Methods of Non-Equilibrium Energy Spectrum (NES) and Differential Non-Equilibrium Energy Spectrum (DNES), Journal of Medicine, Physiology and Biophysics, 20, 13-23.

Ignatov et al., 2015 – Ignatov, I., Mosin, O.V., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, N. (2015). The Evaluation of Mathematical Model of Interaction of Electrochemically Activated Water Solutions (Anolyte and Catholyte) with Water, European Reviews of Chemical Research, 2 (4), 72-86.

Ignatov et al., 2016 – *Ignatov, I., Mosin, O.V., Kirov, P.* (2016). Matematical Model of Kangen Water® Biophysical and Biochemical Effects of Catholyte, *Advances in Physics Theories and Applications*, 51, 33-55.

Ignatov, 2017 – *Ignatov, I.* (2017). Aluminosilicate Mineral Zeolite. Interaction of Water Molecules in Zeolite Table and Mountain Water Sevtopolis from Bulgaria, *Journal of Medicine, Physiology and Biophysics*, 31, 41-45.

Ignatov, Mosin, 2014 – *Ignatov, I., Mosin, O.V.* (2014). Methods for Measurements of Water Spectrum. Differential Non-equilibrium Energy Spectrum Method (DNES), Journal of Health, Medicine and Nursing, 6, 50-72.

Ignatov, Mosin, 2015 – *Ignatov, I., O.V.Mosin.* (2015). Origin of Life and Living Matter in Hot Mineral Water, *Advances in Physics Theories and Applications*, 39, 1-22.

Ignatov, Mosin, 2014a - Ignatov, I., Mosin, O.V. (2014). Mathematical Models of

Distribution of Water Molecules Regarding Energies of Hydrogen Bonds , *Journal of Medicine*, *Physiology and Biophysics*, 6, 50-72.

Ignatov, Mosin, 2014b – *Ignatov, I., Mosin, O.V.* (2014). Structural Models of Water and Ice Regarding the Energy of Hydrogen Bonding, *Nanotechnology Research and Practice*, 7 (3), 96-117.

Ignatov, Mosin, 2014c – *Ignatov*, *I.*, *Mosin*, *O.V.* (2014). Nano Mix of Shungite and Zeolite for Cleaning of Toxins and Increasing of Energy of Hydrogen Bonds among Water Molecules in Human Body, *Journal of Medicine*, *Physiology and Biophysics*, 27, 1-10.

Isaacs et al., 2000 – *Isaacs E.D., Shukla A., Platzman P.M. et al.* (2000). Compton Scattering Evidence for Covalency of the Hydrogen Bonds in Ice, *J. Phys. Chem. Solids*, 61, 403–406.

Krastev et al., 2016 – Krastev, D., Ignatov, I., Mosin, O. V., Penkov, P. (2016). Research on the Effects of the 'Dance of the Spiral' Methodology, with Spectral Analysis of Water Extracts, upon the Physiological Parameters of Plants and the Essential Oil Content, Advances in Physics Theories and Applications, 52, 47-55.

Keutsch, Saykally, 2011 – Keutsch F., Saykally R. (2011). Water Clusters: Untangling the Mysteries of the Liquid, One Molecule at a Time, PNAS, 98(19), 105330–10540.

Krastev et al., 2016 – Krastev, D., Ignatov, I., Mosin, O.V., Penkov, P. (2016). Research of the Effects of the "Dance of the Spiral Metrology" upon the Physiological Parameters of the Plants and the Essential Oil Content, European Journal of Molecular Biotechnology, 11, 29-39.

Kokotkiewicz et al., 2010 – Kokotkiewicz, A., Jaremicz, Z., Luczkiewicz, M. (2010). Aronia Plants: A Review of Traditional Use, Biological Activities, and Perspectives for Modern Medicine, Journal of Medicinal Food, 13 (2), 255-269.

Krasnov, Gordetsov, 2009 – Krasnov, V.V., Gordetsov, A.S. (2009). Infrared Spectral Analysis of Blood Serum as Level of Disturbances of Metabolic Processes in Infusion Children Pathology, Clinical Medicine, 83–94. [in Russian]

Kulling, Rawel, 2008 – Kulling, S.E., Rawel, H.M. (2008). Aronia Melanocarpa – A Review on the Characteristic Components and Potential Health Effects, *Planta Medica*, 74 (13), 1625-34.

Kuriyama et al., 2006 – Kuriyama et al. (2006). Green Tea Consumption and Mortality due to Cardiovascular Disease, Cancer, and all Causes in Japan: The Ohsaki Study, *The Journal of the American Medical Association*, 296 (10), 1255–1265.

Lala et al., 2006 – Lala, G. et al. (2006). Anthocyanin-rich Extracts Inhibit Multiple Biomarkers of Colon Cancer in Rats, Nutr. Cancer, 54, 84-93.

Lambert, Yang, 2003 – Lambert, J. D., Yang, C. S. (2003). Mechanism of Cancer Prevention by Tea Constituents, *Journal of Nutrition*, 133 (10), 3262S-3267S.

Lis et al., 2008 – Lis, G., Wassenaar, L.I., Hendry, M.J. (2008). High-precision Laser Spectroscopy D/H and ¹⁸O/¹⁶O Measurements of Microliter Natural Water Samples. *Anal. Chem.*, 80 (1), 287-293.

Liu et al., 1996 – Liu K., Cruzan J.D., Saykally R.J. (1996). Water Clusters. Science Magazine, 271(5251), 929–933.

Loboda, Goncharuk, 2010 – Loboda O., Goncharuk V. (2010). Theoretical Study on Icosahedral Water Clusters, *Chemical Physics Letters*, 484(4–6), 144–147.

Lis et al., 2008 – Lis, G., Wassenaar, L.I., Hendry, M.J. (2008) High-precision Laser Spectroscopy D/H and ¹⁸O/¹⁶O Measurements of Microliter Natural Water Samples, Anal. Chem., 80(1), 287–293.

Wang et al., 2011 – Lv.J., Wang, W., Krafft, T., Li, Y., Zhang, F., Yuan, F. (2011). Effects of Several Environmental Factors on Longevity and Health of the Human Population of Zhongxiang, Hubei, China, Biol. Trace Elem. Res., 143 (2), 702-716.

Lulli et al., 2012 – Lulli D et al. (2012). Anti-Inflammatory Effects of Concentrated Ethanol Extracts of Edelweiss (Leontopodium alpinum Cass.) Callus Cultures towards Human Keratinocytes and Endothelial Cells, *Hindawi Publishing Corporation Mediators of Inflammation*, 1-12.

Luck et al., 1980 – Luck W., Schiöberg D., Ulrich S. (1980). Infared Investigation of Water Structure in Desalination Membranes. J. Chem. Soc. Faraday Trans., 2(76), 136–147.

Machlin, Bendich, 1988 – *Machlin, L.J., Bendich, A.* (1988). Free Radical Tissue Damage: Protective Role of Antioxidant Nutrients, *The FACEB Journal*, 1(6), 441-445.

Maheshwary et al., 2001 – Maheshwary S., Patel N., Sathyamurthy N., Kulkarni A.D., Gadre S.R. (2001). Structure and Stability of Water Clusters $(H_2O)_n$, n = 8-20, An Ab Initio Investigation. J. Phys. Chem., 105, 10525–10537.

Mariani et al., 2006 – Mariani, E. et al. (2006). Antioxidant Enzyme Activities in Healthy Old Subjects: Influence of Age, Gender and Zinc Status: Results from the Zincage Project, Biogerentology, 7 (5-6), 391-398.

Malhotra, Dhawan, 2008 – Malhotra, A., Dhawan, D.K. (2008). Zinc Improves Antioxidative Enzymes in Red Blood Cells and Hematology in Lithium-Treated rats, Nutr. Res., 28(1), 43-50.

Malik et al., 2003 – Malik, M. et al. (2003). Anthocyanin-rich Extract from Aronia Meloncarpa Induces a Cell Cycle Block in Colon Cancer but not Normal Colonic Cells, Nutr. Cancer, 46,186-196.

Matsumoto et al., 2004 – *Matsumoto, M. et al.* (2004). Gastroprotective Effect of Red Pigments in Black Chokeberry Fruit (Aronia melanocarpa Elliot) on Acute Gastric Hemorrhagic Lesions in Rats, *J. Agric. Food Chem.*, 52 (8), 2226-2229.

Mehandjiev et al., 2017a – Mehandjiev, D., Ignatov, I., Karadzhov, I., Gluhchev, G., Atanasov, A. (2017). On the Mechanism of Water Electrolysis, Journal of Medicine, Physiology and Biophysics, 31, pp. 23-26.

Mehandjiev et al., 2017b – Mehandjiev, D., Ignatov, I., Karadzhov, S., Gluhchev, G., Atanasov, A. (2017.) Processes in Catholyte and Anolyte as Result of Water Electrolysis, European Journal of Molecular Biotechnology, 5 (1), 23-29.

Michaelides, Morgenstern, 2007 – *Michaelides A., Morgenstern K.* (2007). Ice Nanoclusters at Hydrophobic Metal Surfaces, *Nat. Mat.*, 6, 597–599.

Mocchegiani, 2007 – *Mocchegiani, E.* (2007). Zinc, Metallothioneins, and Longevity- Effect of Zinc Supplementation: Zincage Study, *Ann N. Y. Acad. Sci*, 1119, 129-146.

Mosin, Ignatov, 2012a – Mosin, O.V., Ignatov, I. (2012). Isotope Effects of Deuterium in Bacterial and Microalgae Cells at Growth on Heavy Water (D_2O), Voda: Himia i Ecologija, 3, 83–94. [in Russian]

Mosin, Ignatov, 2012b – Mosin, O.V., Ignatov I. (2012). Separation of Heavy Isotopes Deuterium (D), Tritium (T) and Oxygen (¹⁸O) in Water Treatment, Clean Water, Problems and Decisions (Moscow), 3–4. 69-78. [in Russian]

Mosin, Ignatov, 2013 – Mosin, O.V., Ignatov, I. (2013). The Structure and Composition of Natural Carbonaceous Fullerene Containing Mineral Shungite, International Journal of Advanced Scientific and Technical Research, 6(11-12), 9-21.

Pasichnyk, Maggs, 2008 – Pasichnyk I. Everaers R., Maggs A.C. (2008). Simulating van der Waals-interactions in Water/hydrocarbon-based Complex Fluids, J. Phys. Chem. B, 112(6), 1761–1764.

Pauling, 1960 – Pauling L. (ed.) (1960). The Nature of the Chemical Bond and the Structure of Molecules and Crystals, New York: Cornell University Press.

Pimentel, McClellan, 1960 – Pimentel G.C., McClellan A.L. (1960). The hydrogen bond (L. Pouling, Ed), New York: Reinhold Publishing Co.

Pocock, S.J, Shaper, A.G., Packham, R.F. (1981) Studies of Water Quality and Cardiovascular Disease in the United Kingdom, *Sci. Total Environ.*, 18, 25–34.

Olas, 2008 – Olas, B. et al. (2008). Comparative Anti-platelet and Antioxidant Properties of Polyphenol-rich Extracts from Berries of Aronia Melanocarpa, Seeds of Grape and Bark of Yucca Schidigera in vitro, *Platelets*, 19 (1), 70-7.

Orgel, 1963 – *Orgel, L.* (1963). The Maintenance of the Accuracy of Protein Synthesis and its Relevance to Aging, *Biochemistry*, 49, 517–521.

Reisinger et al., 2009 – *Reisinger, U. et al.* (2009). Leogin, the Major Lignin from Edelweiss, Inhibits Intimal Hyperplasia of Venous Bypass Graphs, *Cardiovascular Research*, 82, 524-549.

Ryszawa et al., 2006 – *Ryszawa et al.* (2006). Effects of Novel Plant Antioxidants on Platelet Supreoxide Production and Aggregation Atherosclerosis, *J Physiol Pharmacol*, 57, 611-626

Saykally, 2005 – Saykally, R. (2005). Unified Description of Temperature-dependent Hydrogen Bond Rearrangements in Liquid Water. *PNAS*, 102(40), 14171–14174.

Simonet al., 2006 – Simon, M. et al. (2006). How Much Calcium Is in Your Drinking Water? A Survey of Calcium Concentrations in Bottled and Tap Water and Their Significance for Medical Treatment and Drug Administration, *HSS Journal*, 2(2), 130–135.

Shelton, 1991 – Shelton, R.M. (1991). Aloe Vera: Its Chemical and Therapeutic Properties, Int. Journal Dermatol, 30, 679-683.

Schwalger et al., 2005 – Schwalger, S. et al. (2005). Leontopodic acid—a Novel Highly Substituted Glucaric Acid Derivative from Edelweiss (Leontopodium alpinum Cass.) and its Antioxidative and DNA Protecting Properties, Tetrahedron, 61(19): 4621-4630.

Schwalger et al., 2006 – Schwalger, S. et al. (2006). Development of an HPLC-PAD-MS Assay for the Identification and Quantification of Major Phenolic Edelweiss (*Leontopodium alpium* Cass., *Phytochemical Analysis*, 17 (5), 291-298.

Speroni et al., 2006 – Speroni et al. (2006). In Vivo Efficacy of Different Extracts of Edelweiss (Leontopodium alpinum Cass.) in animal models, Journal of Ethnopharmacology, 89 (2-3): 301-303.

Stewart, 1999 – Stewart, M. et al. (1999). Selenium Compounds have Disparate Abilities to Impose Oxidative Stress and Induce, Apoptosis, Free Radicals Biology and Medicine, 26 (1-2): 42-48.

Sykes, 2007 – *Sykes M.* (2007). Simulations of RNA Base Pairs in a Nanodroplet Reveal Solvation-Dependent Stability. *PNAS*, 104(30), 12336–12340.

Sun et al., 2012 – Sun, Y. et al. (2012). Vitamin E Supplementation Protects Erythrocyte Membranes from Oxidative Stress in Healthy Chinese Middle-aged and Elderly People, Nutrition Research, 32(5), 328-334.

Tokmachev et al., 2010 – Tokmachev A.M., Tchougreeff A.L., Dronskowski R. (2010). Hydrogen-Bond Networks in Water Clusters $(H_2O)_{20}$: An Exhaustive Quantum-Chemical. European Journal of Chemical Physics and Physical Chemistry, 11(2), 384–388.

Tsai C.J., Jordan, 1993 – Tsai C.J., Jordan K.D. (1993). Theoretical Study of the $(H_2O)_6$ Cluster. Chem. Phys. Letters, 213, 181–188.

Valcheva-Kuzmanova, Belcheva, 2006 – Valcheva-Kuzmanova, S.V., Belcheva, A. (2006) Current Knowledge of Aronia Melanocarpa as a Medicinal *Plant. Folia Med.*, 48 (2):11-7.

Valcheva-Kuzmanova et al., 2005 – Valcheva-Kuzmanova, S.V. et al. (2005). Effect of Aronia Melanocarpa Fruit Juice on indomethacin-induced Gastric Mucosal Damage and Oxidative stress in Rats, Experimental and Toxicological Pathology, 56 (6), 385-392.

Vazquez et al., 1996 – *Vazquez, B. et al.* (1996). Anti-inflammatory Activity of Extracts Aloe Vera Gel, J. *Ethnopharmacol*, 55: 69-75.

Vogler, Ernst, 1999 – Vogler, B.K., Ernst. E. (1999). Aloe Vera a Systematic Review of the Clinical Effectiveness, Br. J. Gen: 825-828.

Wang et al., 1997 – Wang H. et al. (1997). Oxygen Radical Absorbing Capacity of Anthocyanins, J. Agric. Food Chem., 45, 304-309.

Wang, Jordan, 2003 – Wang F., Jordan K.D. (2003). A Parallel Tempering Monte Carlo Investigation of $(H_2O)_{6^-}$, *J. Chem. Phys.*, 119, 11645–11653.

Wang et al., 2007 – Wang et al. (2007). Leoligin, the Major Lignan from Edelweiss (*Leontopodium nivale* subsp. *alpinum*), Promotes Cholesterol Efflux from THP-1 Macrophages, *J. Nad. Prod.*, 79(6): 1651–1657.

Woodhead, 1984 – Woodhead, R. (1984). Molecular Biology of Aging, NY, Basic Life Science, 35, pp. 34–37.

Yang et al., 2007 – Yang D.J. et al. (2007). Effects of Different Steeping Methods and Storage on Caffeine, Catechins and Gallic Acid in Bag Tea Infusions, *Journal of Chromatography A*, 1156, 312–320.

Yuan, Shao, 2002 – Yuan, Y., Shao, Y. (2002). Systematic Investigation of Alkali Metal Ion transfer Across the Micro-and-nano-water/1,2-dichloroethane Interfaces Facilated by Dibenzo-18-crown-6, *The Journal of Physical Chemistry B*, 106 (32), 7809–7814.

Zink et al., 2014 – Zink, A. et al. (2014). Outbreak of *Microsporum audouinii* in Munich – the Return of Infectious Fungi in Germany, *Mycose*, 57 (12): 765-770.

Zhao et al., 2004 – *Zhao, C. et al.* (2004). Effects of Commercial Anthocyanin-rich Extracts on Colonic Cancer and Nontumorigenic Colonic Cell Growth. *J Agric Food Chem.*, 52 (20): 6122-8.

Zheng, Wang, 2003a – Zheng, W., Wang, S. Y. (2003). Oxygen Radical Absorbing Capacity of Phenolics in Blueberries, Cranberries, Chokeberries, and Lingonberries, J. Agric. Food Chem., 502-509.

Zheng, Wang, 2003b – Zheng, W., Wang, S.Y. (2003). Oxygen Radical Absorbing Capacity of Phenolics in Blueberries, Cranberries, Chokeberries and Lingonberries, J. Agric Food Chem., 51 (2), 502-509.